arXiv:1603.06444v2 [math.AC] 10 Feb 2017

ON THE SIGN PATTERNS OF THE COEFFICIENTS OF HILBERT POLYNOMIALS

AKIYOSHI TSUCHIYA

Abstract. We show that all sign patterns of the coefficients of Hilbert polynomials of standard graded k-algebras are possible.

The purpose of this paper is to investigate which sign patterns of the coefficients of Hilbert polynomials of standard graded commutative k-algebras are possible. This was asked by Mark Haiman ([2]). In this paper, we give a complete answer to this question.

Let k be a field and R a finitely generated graded k-algebra. We call

$$H(R;i) := \dim_k(R_i)$$

the *Hilbert function* of R, where R_i is the graded component of R of degree i. We say that a k-algebra R is *standard* if it can be finitely generated as a k-algebra by elements of R_1. It is known that if R is standard, then there exist a polynomial $P_R(x) \in \mathbb{Q}[x]$ and a positive integer N such that $H(R;i) = P_R(i)$ for all $i \geq N$ ([3]). We call $P_R(x)$ the *Hilbert polynomial* of a standard graded k-algebra R. We refer the readers to [4] for an introduction to the theory of standard graded k-algebras and Hilbert polynomials.

Let $f(x)$ be a Hilbert polynomial and $d = \deg(f(x))$. The leading coefficient, that is, the coefficient of x^d of $f(x)$ is positive. On the other hand, other coefficients of $f(x)$ are not necessarily positive. We recall that the *sign function* of a real number x is defined as follows:

$$\mathrm{sgn}(x) := \begin{cases} -1 & \text{if } x < 0, \\ 0 & \text{if } x = 0, \\ 1 & \text{if } x > 0. \end{cases}$$

Let $f(x) = \sum_{i=0}^{d} a_i x^i$. For a polynomial $f(x)$, we set

$$S(f(x)) = (\mathrm{sgn}(a_0), \ldots, \mathrm{sgn}(a_{d-1})) \in \{-1, 0, 1\}^d.$$

In this paper, we show that all sign patterns of the coefficients of Hilbert polynomials are possible. In other words, given an integer $d \geq 1$ and $s \in \{-1, 0, 1\}^d$, there exists a Hilbert polynomial $f(x)$ of degree d such that $S(f(x)) = s$. In order to prove

2010 *Mathematics Subject Classification.* 05E40, 13A02.

Key words and phrases. Hilbert polynomial, Hilbert function, standard graded k-algebra.

this question, we consider when a polynomial $f(x) \in \mathbb{Z}[x]$ is a Hilbert polynomial. In fact, we will show the following theorem.

Theorem 1. *Let d be a positive integer and $a_0, \ldots, a_{d-1}$ integers. We set $N = \min\{a_0, \ldots, a_{d-1}\}$. Then the followings hold:*

(1) If $N \geq -2$, then for any integer $n \geq 5d+3$, $nx^d + a_{d-1}x^{d-1} + \cdots + a_0$ is a Hilbert polynomial;

(2) If $N < -2$, then for any integer $n \geq -10d\lfloor N/5 \rfloor + 3$, $nx^d + a_{d-1}x^{d-1} + \cdots + a_0$ is a Hilbert polynomial.

By this theorem, we obtain the following corollary.

Corollary 2. *For any $d \geq 1$ and $s = (s_0, \ldots, s_{d-1}) \in \{-1, 0, 1\}^d$, there exists a Hilbert polynomial $f(x) \in \mathbb{Z}[x]$ of degree d such that $S(f(x)) = s$.*

This corollary says that all sign patterns of the coefficients of Hilbert polynomials are possible.

Before proving Theorem 1, we consider when a monomial $f(x) \in \mathbb{Z}[x]$ is a Hilbert. Now, we collect some results that will be used in the rest of this paper.

Lemma 3 ([1, Theorem 2.5])**.** *Let $f(x) \in \mathbb{Q}[x]$ be such that $f(\mathbb{Z}) \subseteq \mathbb{Z}$, and let $m_0, \ldots, m_d$ be the unique integers such that*

$$f(x) = \sum_{i=0}^{d} \left(\binom{x+i}{i+1} - \binom{x+i-m_i}{i+1} \right),$$

where $d = \deg(f(x))$. Then $f(x)$ is a Hilbert polynomial if and only if $m_0 \geq m_1 \geq \cdots \geq m_d \geq 0$.

This is a numerical characterization of Hilbert polynomials. We call the integers $m_0, \ldots, m_d$ uniquely determined by Lemma 3 the *Macaulay parameters* of $f(x)$, and we write $M(f(x)) = (m_0, \ldots, m_d)$.

Lemma 4 ([1, Theorem 3.5])**.** *Let $f(x), g(x) \in \mathbb{Q}[x]$ be two Hilbert polynomials and let t be a positive integer. Then the following are Hilbert polynomials:*

(i) $f(x) + g(x)$;
(ii) $f(x)g(x)$;
(iii) $tf(x)$;

This lemma gives some fundamental operations on polynomials that preserve the property of being Hilbert polynomials.

By using lemmas, we obtain the following proposition.

Proposition 5. *Let d be a non-negative integer and t a positive integer. We set $f(x) = kx^d$. Then $f(x)$ is a Hilbert polynomial if and only if one of the followings is satisfied:*

(i) $d = 0$;
(ii) $d = 1$ *and* $t \geq 3$;
(iii) $d = 2$ *and* $t \geq 3$;
(iv) $d \geq 3$.

Proof. It is known that when $d = 0$ or $d \geq 3$, x^d is a Hilbert polynomial and x and x^2 are not Hilbert polynomials. (See [1, Theorem 3.8].) By computing the Macaulay parameters of $2x, 3x, 4x, 5x$ and $2x^2, 3x^2, 4x^2, 5x^2$, from Lemma 3 we know $2x, 2x^2$ are not Hilbert polynomials and $3x, 4x, 5x, 3x^2, 4x^2, 5x^2$ are Hilbert polynomials. Hence by Lemma 4, the assertion follows. □

Finally, we prove Theorem 1.

Proof of Theorem 1. It is easy to show that $2x+1$ and $5x-5$ are Hilbert polynomials. Hence, by Lemma 4 and Proposition 5, it follows that $(5x-5)(dx^{d-1}+\cdots+2x+1) = 5(dx^d - x^{d-1} - \cdots - 1)$ is also a Hilbert polynomial.

Assume that $N \geq -2$. Then we know that $5dx^d + a_{d-1}x^{d-1} + \cdots + a_0$ is a Hilbert polynomial. Hence, for any integer $n \geq d+3$, $nx^d + a_{d-1}x^{d-1} + \cdots + a_0$ is a Hilbert polynomial.

Next, assume that $N < -2$. We set $M = -\lfloor N/5 \rfloor$. Then since $-10M \leq 2N < 2N + 3 \leq N$, it follows that $10dMx^d + a_{d-1}x^{d-1} + \cdots + a_0$ is a Hilbert polynomial. Hence for any integer $n \geq 10dM + 3 = -10d\lfloor N/5 \rfloor + 3$, $nx^d + a_{d-1}x^{d-1} + \cdots + a_0$ is a Hilbert polynomial, as desired. □

Acknowledgment. *I would like to thank Professor Jeśus A. De Loera for suggesting me to study the sign patterns of the coefficients of Hilbert polynomials.*

References

[1] F. Brenti, Hilbert polynomials in Combinatorics, *J. Alg. Comb.* **7**(1998), 127–156.
[2] J. A. De Loera, Private communication, 2016.
[3] D. Hilbert, Über die theorie der algebraischen formen, *Math. Annal.* **36**(1890), 473–534.
[4] R. P. Stanley, Combinatorics and Commutative Algebra, Second Ed., Birkhauser, 1996.

(Akiyoshi Tsuchiya) Department of Pure and Applied Mathematics, Graduate School of Information Science and Technology, Osaka University, Suita, Osaka 565-0871, Japan

E-mail address: a-tsuchiya@cr.math.sci.osaka-u.ac.jp